



Emergency Nursing Resource: Family Presence During Invasive Procedures and Resuscitation in the Emergency Department

Does family presence have a positive or negative influence on the patient, family and staff during invasive procedures and resuscitation?

Developed by:

2009 ENA Emergency Nursing Resources Development Committee

Darcy Egging, MS, RN, CEN, CNP
Melanie Crowley, MSN, RN, CEN, MICN
Terri Arruda, MSN, RN, APRN-BC, FNP
Jean Proehl, MN, RN, CEN, CPEN, CCRN, FAEN
Gayle Walker-Cillo, MSN/Ed, RN, CEN, CPEN
AnnMarie Papa, MSN, RN, CEN, NE-BC, FAEN
Suling Li, PhD, RN
Jill Walsh, DNP, RN

ENA Staff Editor

Marlene L. Bokholdt, BSN, RN, CPEN, CCRN-Ped

ENA's Emergency Nursing Resources (ENRs) are developed by ENA members to provide emergency nurses with evidence-based information to utilize and implement in their care of emergency patients and families. Each ENR focuses on a clinical or practice-based issue, and is the result of a review and analysis of current information believed to be reliable. As such, information and recommendations within a particular ENR reflect the current scientific and clinical knowledge at the time of publication, are only current as of their publication date, and are subject to change without notice as advances emerge.

In addition, variations in practice, which take into account the needs of the individual patient and the resources and limitations unique to the institution, may warrant approaches, treatments and/or procedures that differ from the recommendations outlined in the ENRs. Therefore, these recommendations should not be construed as dictating an exclusive course of management, treatment or care, nor does the use of such recommendations guarantee a particular outcome. ENRs are never intended to replace a practitioner's best medical judgment based on the clinical circumstances of a particular patient or patient population. ENRs are published by ENA for educational and informational purposes only, and ENA does not approve or endorse any specific methods, practices, or sources of information. ENA assumes no liability for any injury and/or damage to persons or property arising out of or related to the use of or reliance on any ENR.

Publication Date: December 2009

Table of Contents

	Page
Background/Significance	1
Methodology	1
Evidence Table.....	3
Other Resources Table	3
Summary of Literature Review.....	3
Description of Decision Options/Interventions and the Level of Recommendation	5
Bibliography.....	6
Acknowledgement.....	8

Background/Significance

The concept of family members witnessing the resuscitation or invasive procedure of their relative is one that has garnered much attention over the past few decades. With the rise of family-centered care, family input into healthcare decisions has increased and strict visitation policies have relaxed, even including family at the bedside during invasive procedures and resuscitation. This concept first presented in the early 1980's when Foote Hospital in Michigan began a program to facilitate the practice of family member presence during resuscitation as a response to demands by families (Doyle, 1987). Hanson and Strawser (1992) presented data from that program as the seminal research on this topic. Since then the research has centered on several different aspects of this issue.

Research has been conducted to examine the perspectives of the patient; both children and adults, (Piira, Sugiura, Champion, Donnelly & Cole, 2005; Mortelmans et al., 2009) patients' family members, including opinions regarding family presence and facilitation or hindrance of grief based on witnessing the resuscitation of a relative (Dudley et al., 2009; Mortelmans et al., 2009; Piira et al., 2005; Tinsley et al., 2008) and those of the healthcare team, including opinions regarding family presence and concept family presence being of assistance to or causing interference of the resuscitation or procedure (Basol, Ohman, Simones, & Skillings, 2009; Demir, 2008; Dudley et al., 2009; ENA, 2007; Fallis, McClement & Pereria, 2008; Fernandez et al., 2009; Kuzin et al., 2007; Madden & Condon, 2007; Nigrovic, McQueen, & Neuman, 2007; Piira et al., 2005, Pruitt et al., 2008; Walker, 2008). There are studies that have defined family presence as parental presence for a minor child for invasive procedures (Kuzin, et al., 2007; Nigrovic, McQueen, & Neuman, 2007; Piira, Sugiura, Champion, Donnelly, & Cole, 2005) and resuscitation (Dudley et al., 2009; McGahey-Oakland et al., 2007; Tinsley et al., 2008)) or family members being present during resuscitation of adult relatives (McClement, Fallis, & Pereira, 2009).

Methodology

This ENR was created based on a thorough review and critical analysis of the literature following ENA's [Guidelines for the Development of the Emergency Nursing Resources](#). Via a thorough literature search, all articles relevant to the topic were identified. The following databases were searched: PubMed, eTBLAST, Cochrane - British Medical Journal, Agency for Healthcare Research and Quality (AHRQ; www.ahrq.gov), and the National Guideline Clearinghouse (www.guidelines.gov). Search terms included the key words family presence or parental presence, and invasive procedures, or resuscitation and emergency. Search limitations included articles published in the English language from 2005 to 2009. Systematic, critical and comprehensive reviews included represent earlier works. Classic and seminal research on the issue, as well as non-research articles were also reviewed for historical perspective. In addition, the reference lists of articles found via literature search were scanned for pertinent references.

Articles that met the following criteria were chosen to formulate the ENR: research studies, meta-analyses, systematic reviews, and existing guidelines relevant to the topic. Individuals studies that have been reviewed by any systematic reviews/meta-analyses were not included in the evidence table. Rather, the findings of the systematic reviews/meta-analyses were presented in the evidence table. For example, in 2007, the Emergency Nurses Association published the third edition of *Presenting the Option for Family Presence*. The review of the literature included 117 research studies. Studies in this publication were not individually referenced nor included in the Evidence Table for this ENR. Evidence identified in *Presenting the Option for Family Presence* (3rd ed.) is cited as (ENA, 2007). Other types of articles were also reviewed and provided as additional information. The ENR authors used [standardized](#)

This material is protected by U.S. copyright law. Unauthorized reproduction is prohibited. To request permission to reproduce multiple copies, please e-mail Permissions@ena.org.

[worksheets](#), including Evidence-Appraisal Table Template, Critique Worksheet and AGREE Work Sheet, to prepare tables of evidence ranking each article in terms of the level of evidence, quality of evidence, and relevance and applicability to practice. Clinical findings and levels of recommendations regarding patient management were then made by the Clinical Guidelines Committee according to the ENA's classification of levels of recommendation for practice, which include: Level A High, Level B. Moderate, Level C. Weak or Not recommended for practice (See Table 1).

Table 1. Levels of Recommendation for Practice

<p><u>Level A recommendations: High</u></p> <ul style="list-style-type: none"> • Reflects a high degree of clinical certainty • Based on availability of high quality level I, II and/or III evidence available using Melnyk & Fineout-Overholt grading system (Melnyk & Fineout-Overholt, 2005) • Based on consistent and good quality evidence; has relevance and applicability to emergency nursing practice • Is beneficial
<p><u>Level B recommendations: Moderate</u></p> <ul style="list-style-type: none"> • Reflects moderate clinical certainty • Based on availability of Level III and/or Level IV and V evidence using Melnyk & Fineout-Overholt grading system (Melnyk & Fineout-Overholt, 2005) • There are some minor or inconsistencies in quality evidence; has relevance and applicability to emergency nursing practice • Is likely to be beneficial
<p><u>Level C recommendations: Weak</u></p> <ul style="list-style-type: none"> • Level V, VI and/or VII evidence available using Melnyk & Fineout-Overholt grading system (Melnyk & Fineout-Overholt, 2005) - Based on consensus, usual practice, evidence, case series for studies of treatment or screening, anecdotal evidence and/or opinion • There is limited or low quality patient-oriented evidence; has relevance and applicability to emergency nursing practice • Has limited or unknown effectiveness
<p><u>Not recommended for practice</u></p> <ul style="list-style-type: none"> • No objective evidence or only anecdotal evidence available; or the supportive evidence is from poorly controlled or uncontrolled studies • Other indications for not recommending evidence for practice may include: <ul style="list-style-type: none"> ○ Conflicting evidence ○ Harmfulness has been demonstrated ○ Cost or burden necessary for intervention exceeds anticipated benefit ○ Does not have relevance or applicability to emergency nursing practice • There are certain circumstances in which the recommendations stemming from a body of evidence should not be rated as highly as the individual studies on which they are based. For example: <ul style="list-style-type: none"> ○ Heterogeneity of results ○ Uncertainty about effect magnitude and consequences, ○ Strength of prior beliefs ○ Publication bias

Evidence Table and Other Resources

The articles reviewed to formulate the ENR are described in the [Evidence Table](#). Other articles relevant to family presence were reviewed to serve as additional resources ([Other Resources Table](#)).

Summary of Literature Review

Patient Perspective

There is little evidence to indicate any affect on the patient either experiencing family presence or not. Robinson (as cited in Dingeman, Mitchell, Meyer and Curley, 2007) interviewed three adult survivors of resuscitation and although excluded from the study of family perspectives, all three indicated they were content to have family present. Dudley et al. (2009) reported that parents believed their presence to be helpful to their children during trauma resuscitation. The purpose of their study investigated the effect on resuscitation and testing times, however, they interviewed families for qualitative input. Tinsley et al. (2008) also reported a majority of family members believed their presence during resuscitation was comforting to their child. Mortelmans et al. (2009) interviewed adult inpatients with life-threatening illnesses, asking if they would prefer to have their family members present should their condition deteriorate and they would require resuscitation. A majority indicated that would be their preference despite believing it may be traumatic for them. In a systematic review of literature on parental presence for children undergoing invasive procedures, Piira et al. (2005) reported specifically analyzing pediatric responses. They found mixed results for child behavioral and emotional responses to having parents present. Studies with weaker evidence were more likely to find a significant positive response.

Family Perspective

An increased number of research studies were conducted from the family's perspective. A majority of family members expressed a desire to be present, stating that it is their right to be present and would recommend it to other families (Dingeman et al., 2007; Dudley et al., 2009; McGahey et al., 2007; Mortelmans et al., 2009; Piira et al., 2005; Tinsley et al., 2008). McGahey et al. (2007) identified that all parents surveyed felt the option to be present should be given. Piira et al. (2005) reported that 7 of 17 reviewed studies demonstrated a decreased level of distress and an increased level of satisfaction in families present, while the other 10 studies demonstrated no significant difference between those present and those not. None of the studies reviewed demonstrated any level of increased distress or decreased satisfaction related to being present. Another common theme among family members who had been present during resuscitation is that they believed everything that could have been done for their family member had been done (ENA, 2007; McGahey et al., 2007; Tinsley et al., 2008). Tinsley et al. (2008) also identified a majority of parents (67%) felt it helped them cope with the death of their child, while 43% stated that being present was the thing that helped them the most during the resuscitation.

Health Care Professional Perspective

Results of research that focused on the opinions and perspective of health care professionals indicated support for the practice of family presence during resuscitation and invasive procedures (Basol et al., 2009; Dingeman et al., 2007; ENA, 2007; Fallis, & Pereira, 2009; Kuzin et al., 2007; Madden & Condon, 2007; McClement et al., 2009; O'Connell et al., 2007). Demir (2008) identified that 82% of physicians and

This material is protected by U.S. copyright law. Unauthorized reproduction is prohibited. To request permission to reproduce multiple copies, please e-mail Permissions@ena.org.

nurses in emergency departments and intensive care units in Turkey felt it was inappropriate for families to witness resuscitation. The author suggested that cultural differences may have accounted for the much lower level of approval of the practice in this study compared to others. The author also highlighted that 86% of the respondents had read no research or other articles regarding family presence and 97% had no knowledge of international guidelines published on the topic.

Common themes emerged from many of the studies regarding the opinions of health professionals. Those approving of family presence thought it helped them to see the effort of the resuscitation team; that everything that could have been done, had been done, which may lower the risk of litigation surrounding the resuscitation or procedure (Basol et al., 2009; Critchell & Marik, 2007; Dingeman et al., 2007; McClement et al., 2009; Pruitt et al., 2008; Walker, 2008). Another theme was that health care professionals felt family presence was a positive experience; that it humanized the patient and supported patient dignity (Basol et al., 2009; Critchell & Marik, 2007; Demir, 2008; McClement et al., 2009; Pruitt et al., 2008). Many studies demonstrated that health care professionals felt having family members present enhanced communication and facilitated education (Basol et al., 2009; Dingeman et al., 2007; Kuzin et al., 2007; McClement et al., 2009; Pruitt et al., 2008; Walker, 2008). Another theme from health care professionals was that it facilitated the grief process in the case of unsuccessful resuscitation. It gave family members the opportunity to say good-bye and promoted families' acceptance of the death of their loved one (Demir, 2008; Dingeman et al., 2007; McClement et al., 2009; Walker, 2008).

Despite the majority of health care professionals having expressed support for the concept of family member presence during resuscitation and procedures there were several themes that emerged demonstrating reasons for reservation regarding the practice. These themes included the possibility of families interfering with the process and disrupting care (Basol et al., 2009; Demir, 2008; Dingeman et al., 2007; Fernandez et al., 2009; Madden & Condon, 2007; McClement et al., 2009; Walker, 2008), increased performance anxiety and stress on the part of clinicians, the interference with the process of teaching (Basol et al., 2009; Demir, 2008; Dingeman et al., 2007; Fernandez et al., 2009; Madden & Condon, 2007; McClement et al., 2009; Walker, 2008), the possibility that witnessing the event may be too traumatic for families (Basol et al., 2009; Demir, 2008; Dingeman et al., 2007; Fernandez et al., 2009; McClement et al., 2009; Pruitt et al., 2008; Walker, 2008), and misinterpretation of procedure and increased risk of litigation related to families witnessing resuscitation and procedures (Demir, 2008; Dingeman et al., 2007; Fernandez et al., 2009; Madden & Condon, 2007; McClement et al., 2009; Walker, 2008).

Interference with Care

One final group of studies observed the care provided during resuscitations and procedures to determine any demonstrable effect on performance by health care professionals. O'Connell et al. (2007) researched pediatric trauma activations and identified no significant difference in time to log-rolling, radiographs, intravenous access, central line placement, intubation or chest tube insertion based on family members having been present in the trauma room. The authors reported no interference of care by any family member in the 196 cases included in the study that had family members present. Dudley et al. (2009) also examined pediatric trauma resuscitations in the cases of 705 patients and discovered no significant delay in time to computerized tomography or change in resuscitation times for patients with family members present in the trauma room. Nigrovic et al. (2009) examined success rates for lumbar puncture of over 1400 pediatric patients and found no significant correlation between family member presence and traumatic or unobtainable lumbar punctures. Sacchetti et al. (2005) observed 37

This material is protected by U.S. copyright law. Unauthorized reproduction is prohibited. To request permission to reproduce multiple copies, please e-mail Permissions@ena.org.

pediatric patients undergoing invasive procedures and reported 2 cases to have had minor interruptions by present family members. Both procedures continued after adequate education of the family member with no significant delay of care. Basol et al. (2009) reported on the institution of a policy providing family members the option to be present during resuscitation of a family member and the authors stated no interference of care or negative experiences with family members. The single study that indicated interference with performance was done by Fernandez et al. (2009) with 2nd and 3rd year emergency medicine residents in the simulation laboratory performing resuscitation scenarios. This study demonstrated a significant delay to initiation of cardiopulmonary resuscitation and medication administration in those groups with simulated family members present. The author stated that although the simulation used in this study was high-fidelity, it was still simulation.

Family Member Presence Policy

Basol et al. (2009) relate their experiences with implementing a policy regarding family member presence. The staff stated support for the concept, but moreover stated the need for a policy to provide consistency, guidelines and improve communication among the team. Madden & Condon (2007) surveyed 90 trauma nurses in Ireland and found that 74% preferred to have a written policy on the practice, although none had one in place.

Description of Decision Options/Interventions and the Level of Recommendation

Conclusions and recommendations about family presence during invasive procedures and resuscitation in the emergency department:

- There is some evidence that patients would prefer to have their family members present during resuscitation.
- There is strong evidence that family members wish to be offered the option to be present during invasive procedures and resuscitation of a family member.
- There is little or no evidence to indicate that the practice of family member presence is detrimental to the patient, the family or the health care team.
- There is evidence that family member presence does not interfere with patient care during invasive procedures or resuscitation.
- There is evidence that health care professionals support the presence of a designated health care professional assigned to present family members to provide explanation and comfort.
- There is some evidence that a policy regarding family member presence provides structure and support to health care professionals involved in this practice.
- Family member presence during invasive procedures or resuscitation should be offered as an option to appropriate family members and should be based on written institution policy (Level B).

Bibliography

- Agard, M. (2008). Creating advocates for family presence during resuscitation. *Medsurg Nursing, 17*(3), 155-160.
- Basol, R., Ohman, K., Simones, J., & Skillings, K. (2009). Using research to determine support for a policy on family presence during resuscitation. *Dimensions in Critical Care Nursing, 28*(5), 237-247.
- Critchell, C.D. & Marik, P.E. (2007). Should family members be present during cardiopulmonary resuscitation? A review of literature. *American Journal of Hospice & Palliative Care, 24*(4), 311-317.
- Demir, F. (2008). Presence of patients' families during cardiopulmonary resuscitation: Physicians' and nurses' opinions. *Journal of Advanced Nursing, 63*(4), 409-416.
- Dingeman, R.S., Mitchell, E.A., Meyer, E.C., & Curley, M.A. (2007). Parent presence during complex invasive procedures and cardiopulmonary resuscitation: A systematic review of literature. *Pediatrics, 120*(4), 842-854.
- Doyle, C.J., Post, H., Burney, R.E., Maino, J., Keefe, M., & Rhee, K.J. (1987). Family participation during resuscitation: An option. *Annals of Emergency Medicine, 16*(6), 673-675.
- Dudley, N.C., Hansen, K.W., Furnival, R.A., Donaldson, A.E., Van Wagene, K.L., & Saife, E.R. (2009). The effect of family presence on the efficiency of pediatric trauma resuscitations. *Annals of Emergency Medicine, 53*(6), 777-784e3.
- Emergency Nurses Association. (2007). Presenting the option for family presence. (3rd ed.). Des Plaines, IL: Author. Chapter 2: "Review of the Literature on Family Presence".
- Fallis, W.M., McClement, S., Pereira, A. (2008). Family presence during resuscitation: A survey of Canadian critical care nurses' practices and perceptions. *Dynamics, 19*(3), 22-28.
- Farah, M.M., Thomas, C.A., Shaw, K.N. (2007, August). Evidence-based guidelines for family presence in the resuscitation room: A step-by-step approach. *Pediatric Emergency Care, 23*(8), 587-591.
- Fernandez, R., Compton, S., Jones, K.A., & Velilla, M.A. (2009). The presence of a family witness impacts physician performance during simulated medical codes. *Critical Care Medicine, 37*(6), 1956-1960.
- Fineout-Overholt, E., Melynk, B., & Schultz, H. (2005). Transforming health care from the inside out: Advancing evidence-based practice in the 21st Century; *Journal of Professional Nursing 21*(6), 335-344.
- Hanson, C., & Strawser, D. (1992). Family presence during cardiopulmonary resuscitation: Foote Hospital emergency department's nine-year perspective. *Journal of Emergency Nursing, 18*(2), 104-106.
- Kuzin, J.K., Yborra, J.G., Taylor, M.D., Chang, A.C., Altman, C.A., Whitney, G.M., & Mott, A.R. (2007). Family-member presence during interventions in the intensive care unit: Perceptions of pediatric cardiac intensive care providers. *Pediatrics, 120*(4), e895-e901.
- Madden & Condon, C. (2007). Emergency nurses' current practices and understanding of family presence during CPR. *Journal of Emergency Nursing, 33*(5), 433-440.
- McClement, S.E., Fallis, W.M., & Pereira, A. (2009). Family presence during resuscitation: Canadian critical care nurses' perspectives. *Journal of Nursing Scholarship, 41*(3), 233-240.
- McGahey-Oakland, P.R., Lieder, H.S., Young, A. & Jefferson, L.S. (2007). Family experiences during resuscitation at a children's hospital emergency department. *Journal of Pediatric Health Care, 21*(4), 217-225.
- Mortelmans, L.J.M., Van Broeckhoven, V., Van Boxtael, S., De Cauwer, H.G., Verfaillie, L., Van Hellemond, P.L.A., et al. (2009). Patients' and relatives' view on witnessed resuscitation in the emergency department: A prospective study. *European Journal of Emergency Medicine, 00*, 000-000. (Ahead-of-print publication).
- Nigrovic, L.E., McQueen, A.A., Neuman, M.I. (2007). Lumbar success rate is not influenced by family-member presence. *Pediatrics, 120*(4), e777-e782.
- O'Connell, K.J., Farah, M.M., Spandorger, P., & Zorc, J.J. (2007). Family presence during pediatric trauma
- This material is protected by U.S. copyright law. Unauthorized reproduction is prohibited. To request permission to reproduce multiple copies, please e-mail Permissions@ena.org.

- team activation: An assessment of a structured program. *Pediatrics*, 120(3), e565-e574.
- Piira, T., Sugiura, T., Champion, G.D., Donnelly, N., & Cole, A.S.J. (2005). The role of parental presence in the context of children's medical procedures: A systematic review. *Child: Care, Health & Development*, 31(2), 233-243.
- Pruitt, L.M., Johnson, A., Elliott, J.C., & Polley, K. (2008). Parental presence during pediatric invasive procedures. *Journal of Pediatric Health Care*, 22(2), 120-127.
- Sacchetti, A., Paston, C., & Carraccio, C. (2005). Family members do not disrupt care when present during invasive procedures. *Academic Emergency Medicine*, 12(5), 477-479.
- Tinsley, C., Hill, J.B., Shar, J., Zimmerman, G., Wilson, M., Freier, K., & Abd-Allah, S. (2008). Experience of families during cardiopulmonary resuscitation in a pediatric intensive care unit. *Pediatrics*, 122(4), e799-e804.
- Walker, W. (2007). Accident and emergency staff opinion of the effects of family presence during adult resuscitation: critical literature review. *Journal of Advanced Nursing*, 61(4), 348-362.

© Emergency Nurses Association, 2009.

Acknowledgement

ENA would like to acknowledge the 2009 Institute for Emergency Nursing Research (IENR) Committee for the review of this document. Members of the IENR Committees include:

Cynthia L. Dakin, PhD, RN, Chairperson
Gail Pisarcik Lenehan, EdD, RN, FAEN, FAAN, ENA Board of Directors Liaison
Sarah Anderson, PhD, RN, CEN, SANE-A
Susan Barnason, PhD, RN, APRN-CNS, CEN, CCRN
Gordon Gillespie, PhD, RN, PHCNS-BC, CEN, CPEN, CCRN, FAEN
Vicki Keough, PhD, RN, ACNP
Stephen Stapleton, PhD(c), RN, CEN

ENA also acknowledges the assistance of Leslie Gates in coordinating the work of the Committee.